

Food and Drug Administration, HHS

§ 522.778

suitable dosage level depending upon the species.

(3) For use in wild or exotic animals only. Do not use in domestic food-producing animals. Do not use 30 days before, or during, the hunting season in free-ranging wild animals that might be used for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian. Distribution is restricted to veterinarians engaged in zoo and exotic animal practice, wildlife management programs and researchers.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 16241, Apr. 15, 1983; 60 FR 39847, Aug. 4, 1995; 64 FR 15684, Apr. 1, 1999]

§ 522.770 Doramectin.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of doramectin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.225 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Amount.* 200 micrograms per kilogram (10 milligrams per 110 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, sucking lice, and mange mites. To control infections and to protect from reinfection with *Cooperia oncophora* and *Haemonchus placei* for 14 days, *Ostertagia ostertagi* for 21 days, and *C. punctata*, *Oesophagostomum radiatum*, and *Dictyocaulus viviparus* for 28 days after treatment.

(iii) *Limitations.* Administer as a single subcutaneous or intramuscular injection. Do not slaughter cattle within 35 days of treatment. Not for use in female dairy cattle 20 months of age or older. Do not use in calves to be processed for veal.

(2) *Swine*—(i) *Amount.* 300 micrograms per kilogram (10 milligrams per 75 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, kidney worms, sucking lice, and mange mites.

(iii) *Limitations.* Administer as a single intramuscular injection. Do not slaughter swine within 24 days of treatment. Consult your veterinarian for as-

sistance in the diagnosis, treatment, and control of parasitism.

[61 FR 53321, Oct. 11, 1996, as amended at 62 FR 44410, Aug. 21, 1997; 62 FR 62242, Nov. 21, 1997; 63 FR 68183, Dec. 10, 1998; 64 FR 13509, Mar. 19, 1999]

§ 522.775 Doxapram hydrochloride injection.

(a) *Specifications.* The drug is a sterile aqueous solution containing 20 milligrams doxapram hydrochloride per milliliter.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) Administer to dogs, cats, and horses to stimulate respiration during and after general anesthesia; to speed awakening and return of reflexes after anesthesia. Administer to neonate dogs and cats to initiate respiration following dystocia or caesarean section; to stimulate respiration following dystocia or caesarean section.

(2) For intravenous use in dogs and cats at a dose of 2½ to 5 milligrams of doxapram hydrochloride per pound of body weight in barbiturate anesthesia, 0.5 mg per lb. in gas anesthesia; for intravenous use in horses at 0.25 mg per lb. of body weight in barbiturate anesthesia, 0.2 mg per lb. in inhalation anesthesia, 0.25 mg per lb. with chloral hydrate with or without magnesium sulfate; for subcutaneous, sublingual, or umbilical vein administration in neonate puppies at a dose rate of 1 to 5 mg; for subcutaneous or sublingual use in neonate kittens at 1 to 2 mg. Dosage may be repeated in 15 to 20 minutes if necessary.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 17838, Apr. 23, 1975, as amended at 67 FR 67521, Nov. 6, 2002]

§ 522.778 Doxycycline hyclate.

(a) *Specifications.* Doxycycline hyclate solution contains 8.5 percent doxycycline activity. A syringe of *N*-methyl-2-pyrrolidone and poly (DL-lactide) mixed with a syringe of doxycycline produces 0.5 milliliter of solution.

(b) *Sponsor.* See 000009 in § 510.600(c) of this chapter.

(c) [Reserved]